

Research Article

Medicinal plant *Potentilla fulgens* and its effect *in vitro* against *Fasciola gigantica*

Pradeep Kumar*

Department of Zoology, S.G.N. Government P.G College, Muhammadabad Gohna, Mau-276403, UP, India

Abstract

Fascioliasis is a one of the most important serious parasitic zoonotic disease which caused by trematode giant liver fluke *Fasciola hepatica* and *F. gigantica* among cattle's and humans. The infection of *Fasciola* can be control by the use of phytochemicals as anthelmintic components. The anthelmintic activities of dried root powder of medicinal plant *Potentilla fulgens* and their different preparations (organic extracts and column purified fraction) are uses *in vitro* against liver fluke *F. gigantica*. The dried root powder, different organic extract, and column fractions were time and concentration-dependent. Among all the organic extracts, ethanol extract was high toxic than other organic extracts. The toxic effect of ethanolic extract of *P. fulgens* after 2h exposure the LC₅₀ value is 5.22 mg/ml against *F. gigantica*. The column purified fraction of dried root powder of *P. fulgens* shows more toxicity. The 2h LC₅₀ of column purified fraction was 3.25 mg/ml whereas in 8h exposure the LC₅₀ is 1.24 mg/ml. The phytochemicals of the *P. fulgens* may be used as anthelmintic components against liver fluke *F. gigantica*.

Introduction

The liver fluke *Fasciola hepatica* and *F. gigantica* is the causative agent of fasciolosis in cattle and human populations [1]. These diseases are parasitic in livestock with over 700 million production animals at risk of infection [2]. Fascioliasis is mainly related to plant-borne trematode digenetic zoonotic disease. The definite host of liver fluke is cattle, sheep, buffaloes, and goats which have an impact on the development, growth rate, the productivity of the ruminants and it's considered economically significant [3,4]. Fasciolosis is very common in the cattle population of the eastern part of Uttar Pradesh, India [5-10]. In humans, the adult fluke of *Fasciola* causes a variety of symptoms such as long-standing fever, malaise, weight loss, eosinophilia, pain under the right costal margin, and anemia due to feeding on the blood [11].

Anthelmintic synthetic drugs can be used for the control of liver fluke infection, but it causes adverse effect on the host as well as develops resistant/residual effect in the fluke. Synthetic drugs are not easily available in some of the remote rural areas of developing countries [12]. The plant products or active phytochemicals are may be an effective anthelmintic component. Therefore, the use of the common medicinal plant as anthelmintic offers an alternative source that can solve these

problems and it may be more acceptable due to eco-friendly and easily available for the users [13]. The medicinal plant *Potentilla fulgens* are commonly used as antihyperglycemic, antioxidant, antitumor, anti-hyperlipidemic, antiulcerogenic, anti-inflammatory, anthelmintic larvicidal, and molluscicidal [14-17]. The present study aims to evaluate the different preparations of medicinal plant *P. fulgens in vitro* treatments and their anthelmintic efficacy against the giant liver fluke *F. gigantica*.

Material and methods

Collection of liver fluke (*F. gigantica*)

The adult fluke *F. gigantica* (3.2 ± 0.18 cm in length) were collected from the infected bile ducts of the freshly slaughtered buffaloes from slaughterhouse district Gorakhpur (UP), India. The live fluke was kept in freshly prepared Hedon-Fleig (H-F) solution at 37 ± 2 °C in a BOD incubator until use.

Plants

The dried root of the *Potentilla fulgens* was procured from the local market in Muhammadabad Gohna, district Mau (UP), India, and authenticity by Dr. A. K. Singh, Department of Botany, S.G.N. Government P.G. College, Muhammadabad Gohna, Mau, Uttar Pradesh.

More Information

*Address for Correspondence: Pradeep Kumar, Department of Zoology, S.G.N. Government P.G College, Muhammadabad Gohna, Mau-276403, UP, India, Email: pkumar_gpu@yahoo.co.in

Submitted: May 03, 2021

Approved: May 12, 2021

Published: May 13, 2021

How to cite this article: Kumar P. Medicinal plant *Potentilla fulgens* and its effect *in vitro* against *Fasciola gigantica*. Insights Clin Cell Immunol. 2021; 5: 004-007.

DOI: 10.29328/journal.icci.1001018

Copyright: © 2021 Kumar P. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Keywords: *Potentilla fulgens*; *Fasciola gigantica*; Fascioliasis; Anthelmintic activity



Preparation of crude plant products

The freshly dried root of *P. fulgens* washes through fresh-water and dry under sunlight till to dry and cut into small pieces. The dried root pieces of *P. fulgens* were pulverized in the electric grinder machine and the crude powders thus obtained were used for the *in vitro* anthelmintic activity against *F. gigantica*.

Organic solvent extracts

Five gram dried powder of *P. fulgens* were extracted with 500 ml each of 98.1% acetone, 97.8% ether, 98.6% chloroform, and 94.7% ethanol at room temperature for 48h. The solvents were removed under a vacuum machine and the following remaining dried parts were used for the determination of *in vitro* anthelmintic activity of *P. fulgens* and extract obtained 350 mg-ethanol, 360 mg-chloroform, 385 mg-ether, and 395 mg-acetone.

Column purified-fractions

One thousand milliliters of 94.7% ethanol were used for the fractions of dried root powder of *P. fulgens* were subjected to silica gel through 5 × 45 cm column chromatography (60-120 mesh, Qualigens Glass, Purchases from Precious Electrochemidus Private Limited, Bombay, India) for different fractions. One hundred milliliter fractions eluted with ethanol were collected. Ethanol from the column purified fractions was evaporated under a vacuum machine and the remaining solids column extracts were used for the determination of *in vitro* anthelmintic activity against *Fasciola*.

Hedon-Fleig (H-F) solution

The Hedon-Fleig (H-F) solution is used in this study which was prepared by the method of Hajare, et al. [18]. H-F solution contain NaCl-119.82 mM, MgSO₄-0.29 mM, KCl-4.01 mM, CaCl₂-0.40 mM, NaHCO₃-17.8 mM, Glucose-22.3 mM, Streptomycin sulphate-6900 unit 10 mg/l and Benzyl penicillin-9900 units/l at 38 ± 2 °C in BOD incubator.

In vitro toxicity determination

In vitro toxicity of dried root powder, organic extract (acetone, ether, chloroform, methanol, and ethanol), and column purified-fraction of *P. fulgens* were performed in the petri dish by the method of Kumar and Singh, [19]. Six Petri dishes were set for each concentration of different preparations of the *P. fulgens* against *F. gigantica*. Ten *F. gigantica* were kept in each Petri dish (10 cm × 1.5 cm) containing 50 ml H-F solution. Flukes were exposed to different concentrations of the different preparations (Table 1). The mortality of adult *Fasciola* was observed after 2h, 4h, 6h, and 8h exposure. The control group of the experiment was kept in an equal volume of H-F solution in a Petri dish under similar laboratory conditions but without treatment. The mortality of *Fasciola* was established by the opening of the sucker and contraction of the body. Usually in H-F solution in *in vitro* *F. gigantica* can

survive up to 48h. Mortality of fluke was observed after 2h up to 8h were counted in treated and control groups.

In vitro, anthelmintic toxicity data were observed every 2h up to 8h. The Lethal Concentration (LC₅₀) values, lower and upper confidence limits (LCL and UCL), slope values, and t-ratio value were calculated by the POLO computer program [20].

Results

In vitro anthelmintic activity of dried root powder of *P. fulgens* and their different preparations against *F. gigantica* was concentration and time-dependent. The lethal concentration (LC₅₀) values of dried root powder at 2h, 4h, 6h, and 8h were 8.35, 7.12, 6.45, and 5.78 mg/ml, respectively (Table 1). Whereas, among all the organic extract (acetone, ether, chloroform, methanol, and ethanol) the ethanol extract were more toxic. The 2h, 4h, 6h, and 8h LC₅₀ of ethanol extract of dried root powder of *P. fulgens* against *F. gigantica* were 5.22, 5.02, 4.88, and 3.43 mg/ml, respectively. The column purified fractions of dried root powder of *P. fulgens* were highly toxic. The 2h, 4h, 6h, and 8h exposure the LC₅₀ value of the column purified-fractions were observed 3.25, 2.65, 1.94, and 1.24 mg/ml, respectively (Table 1). The slope values given in table 1 were steep and the separate estimates of LC₅₀ values based on each of the six replicates of the experiments were found to be within the 95% confidence limits of lethal concentrations. The t-ratio value is greater than 1.96 that indicates the significant anthelmintic efficacy of the various treatments (Table 1).

Discussion

The present study is demonstrated that the dried root powder, organic extracts, and column purified fractions of *P. fulgens* are potent sources of anthelmintic components against *F. gigantica*. It may be possible that active phytochemicals of the medicinal plant, *P. fulgens* are entering the body of fluke *F. gigantica* and cause mortality. In the treated group, all the preparations of the *P. fulgens* are cause significant toxicity *in vitro* against the *F. gigantica* (Table 1). But no mortality was observed in the control group. It indicates that the active phytochemicals of *P. fulgens* are entering through the tegument layer which caused paralysis and mortality of the fluke. The toxicity of different preparations of *P. fulgens* is time as well as concentration dependant as evident from the LC₅₀ values and exposure period (Table 1). Higher toxicity of ethanol extract was observed among all organic extracts which indicate that the anthelmintic components *P. fulgens* are more soluble in ethanol organic solvent.

In vitro toxicity of the root powder of the *P. fulgens* and their different extract preparations are significant and cause antilarvicidal activities against sporocyst, redia, and cercaria larva of the *F. gigantica* [21]. In *in vitro* treatment at 2h exposure, the highest toxicity was noted against sporocyst,

Table 1: *In vitro* anthelmintic activity of dried root powder of *P. fulgens* and their different organic extract, column purified against *F. gigantica* at different exposure periods.

Exposure periods	Values	<i>In vitro</i> exposure of Helminthicides (mg/ml)						
		<i>Potentilla fulgens</i> dried root powder	Acetone extract	Ether extract	Chloroform extract	Methanol extract	Ethanol extract	Column purified
2h	LC ₅₀	8.35	7.76	7.21	6.45	6.95	5.22	3.25
	LCL	6.42	6.12	6.58	5.45	5.72	4.25	2.12
	UCL	9.24	8.45	8.93	7.82	8.64	7.93	4.65
	Slope-value	1.30 ± 0.21	1.12 ± 0.20	1.27 ± 0.11	1.18 ± 0.19	1.23 ± 0.27	1.19 ± 0.17	1.25 ± 0.16
	t-ratio	3.45	4.32	3.55	4.11	3.23	3.96	4.12
4h	LC ₅₀	7.12	6.45	6.37	5.86	6.01	5.02	2.65
	LCL	6.89	5.35	5.37	4.61	4.65	4.12	1.75
	UCL	8.23	7.91	8.32	6.94	7.74	7.53	3.24
	Slope-value	1.14 ± 0.18	1.15 ± 0.21	1.26 ± 0.29	1.27 ± 0.11	1.19 ± 0.24	1.11 ± 0.23	1.31 ± 0.25
	t-ratio	3.11	4.43	3.76	4.26	3.76	3.23	4.23
6h	LC ₅₀	6.45	5.54	6.12	5.12	5.76	4.88	1.94
	LCL	5.96	4.55	5.13	4.55	4.71	3.24	0.66
	UCL	7.12	6.35	8.93	7.32	6.77	5.35	2.13
	Slope-value	1.35 ± 0.20	1.33 ± 0.19	1.27 ± 0.22	1.18 ± 0.35	1.26 ± 0.16	1.39 ± 0.26	1.35 ± 0.21
	t-ratio	4.54	3.14	4.54	3.23	3.65	4.87	4.53
8h	LC ₅₀	5.78	4.87	5.86	4.95	4.64	3.43	1.24
	LCL	4.86	3.24	4.32	3.91	3.74	2.46	0.56
	UCL	6.31	6.51	7.43	5.88	6.27	4.99	2.02
	Slope-value	1.26 ± 0.19	1.22 ± 0.24	1.26 ± 0.17	1.34 ± 0.29	1.17 ± 0.24	1.34 ± 0.26	1.32 ± 0.26
	t-ratio	3.57	4.56	3.11	4.12	4.23	3.54	4.13

Ten *Fasciola gigantica* in six batches were exposed *in vitro* (H-F solution) on different concentrations of the above anthelmintic preparations. Mortality of *Fasciola* was determined after every 2 h exposure period. LCL: Lower Confidence Limits; UCL: Upper Confidence Limits

redia, and cercaria the LC₅₀ value of column extract were 62.42, 59.25, and 45.11 mg/ml, respectively. It may be due to the uptake of the active moiety which progressively increases in the fluke body with an increase in the exposure period. It may be possible that the phytochemicals of the *P. fulgens* in the *F. gigantica* could change the different enzyme activity and cause effects. Many numbers of medicinal plants are have been used for the treatment of parasitic infection in animals and humans [22]. Kumar and Singh, [19] have been reported that in *in vivo* the common species *Allium sativum*, *Ferula asafoetida*, *Syzygium aromaticum* and their active components like allicin, ferulic acid, umbelliferone, and eugenol have anthelmintic activities against *F. gigantica*. The binary combinations (1:1 ratio) of the active components the allicin, ferulic acid, umbelliferone, and eugenol are more effective *in vitro* against the liver fluke *F. gigantica* [23]. The ethanolic root extract of *P. fulgens* is preventing gastric ulcers in rats due to H⁺ K⁺-ATPase inhibitory and antihistaminic activities [24]. The dried root powder of *P. fulgens* and their different organic extracts, column purified fractions are a potent source of the molluscicides which cause significant mortality against liver fluke vector snails *Lymnaea acuminata* and *Indoplanorbis exustus* [15-17]. Roy, et al. [25] have been reported that the alcoholic extract of dried root powder of *P. fulgens* has significantly reduced the vital tegumental enzymatic activity of the alkaline phosphatase, acid phosphatase, and adenosine triphosphatase (ATPase) in cestodes parasite *Raillietina echinobothrida* and trematodes *Gastrothylax crumenifer*. The root extract of *P. fulgens* has flavonoids and tannin in high amounts [26]. *In vitro* and *in vivo* studies have been evidence to support the anthelmintic effect which feeds the tannins and other polyphenols against abdominal and intestinal

parasitic nematodes [27,28]. Athanasiadou, et al. [29] have been evaluated that tannin shows anthelmintic activities against infected sheep with *Trichostrongylus colubriformis* and causing larval mortalities. The compound of tannin in *in vitro* inactivates the enzyme activities in *Trichostrongylus colubriformis* larvae which are responsible for hatching and development [30]. Previously in different studies have been evaluated that the root extract of *P. fulgens* possesses antitumor [31], antioxidant [26], anthelmintic [25], and gastroprotective [24]. The root extract of *P. fulgens* in *in vitro* inhibit enzyme activities of amylase, α -glucosidase, β -glucosidase, and lipase in the liver, kidney, and eye lens of the diabetic mice [32].

Conclusion

It can be concluded from the present study that the dried root powder of *P. fulgens* and their different organic extract and column purified fractions may be used as potent anthelmintic components for the control of liver fluke *F. gigantica*. This study also revealed a further study that the phytochemicals of the *P. fulgens* at the molecular level elucidate in the parasitic life of the liver fluke *F. gigantica*.

References

- Phalee A, Wongsawad C, Rojanapaibul A, Chai J. Experimental life history and biological characteristics of *Fasciola gigantica* (Digenea: Fasciolidae). Korean J Parasitol. 2015; 53: 59-64. PubMed: <https://pubmed.ncbi.nlm.nih.gov/25748710/>
- Chen JX, Chen MX, Ai L, Xu XN, Jiao JM. An Outbreak of Human Fascioliasis *gigantica* in Southwest China. PLoS ONE. 2013; 8: 71520. PubMed: <https://pubmed.ncbi.nlm.nih.gov/23951181/>
- Kuchai JA, Chisnti MZ, Zaki MM, Rasool SAM, Ahmad J, et al. Some epidemiological aspects of fascioliasis among cattle of Ladakh. Global Vet. 2011; 7: 342-346.

4. Eshetu EN, Thomas A, Awukew A, Goa, Buyako B. Study on the prevalence of Bovine Fascioliasis and Estimated financial loss due to liver condemnation: In case of Angacha Woreda, Kambata Tembaro Zone, Southern Ethiopia. *J Biol Agric Healthcare*. 2017; 7: 78-83.
5. Singh O, Agarwal RA. Toxicity of certain pesticides to two economic species of snails in northern India. *J Economic Entomol*. 1981; 74: 568-571.
6. Kumar P, Singh DK. Molluscicidal activity of *Ferula asafoetida*, *Syzygium aromaticum* and *Carum carvi* and their active components against the snail *Lymnaea acuminata*. *Chemosphere*. 2006; 63: 1568-1574. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/16310827/>
7. Singh V, Singh DK. The effect of abiotic factors on the toxicity of cypermethrin against the snail *Lymnaea acuminata* in the control of fascioliasis. *J Helminthol*. 2009; 83: 39-45.
8. Kumar P, Singh VK, Singh DK. Enzyme activity in the nervous tissue of *Lymnaea acuminata* fed to different bait formulations. *Am J Chem*. 2012; 2: 89-93.
9. Singh RN, Kumar P, Kumar N, Singh DK. Efficacy of binary combination of deltamethrin+MGK-264 on levels of biochemical changes in the snail *Lymnaea acuminata*. *Int J Pharm Pharmaceut Sci*. 2020; 12: 111-116.
10. Vishwakarma AK, Kumar P. In vivo Anthelmintic Activity of Medicinal plant *Asparagus racemosus* against Larva of *Fasciola gigantica*. *Res J Agri Sci*. 2021; 12: 675-680.
11. Ghoneim MA, El-kirdasy AF, Goher AK, Moussa WM, El-Ballal SS, et al. Expression of cathepsin L1 Gene of *Fasciola gigantica* in different development stages. *J Appl Sci Res*. 2008; 4: 2017-2078.
12. Singh D, Swarnkar CP, Khan FA. Anthelmintic resistance in gastrointestinal nematodes of livestock in India. *J Vet Parasitol*. 2002; 16: 115-130.
13. Hammond JA, Fielding D, Bishop SC. Prospects for plant Anthelmintics in tropical veterinary medicine. *Vet Res Com*. 1997; 21: 213-228. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/9090048/>
14. Kaul K, Jaitak V, Kaul VK. Review on pharmaceutical properties and conservation measures of *Potentilla fulgens* Wall. Ex Hook-a medicinal endangered herb of higher Himalaya. *Indian J Nat Prod Resour*. 2011; 2: 298-306.
15. Kumar P, Sunita K, Singh DK. Efficacy of *Potentilla fulgens* root powder and their different organic extract against fresh water vector snail *Lymnaea acuminata*. *Asian J Animal Vet Adv*. 2018a; 13: 3034.
16. Kumar P, Sunita K, Singh DK. Molluscicidal activity of different organic root extract of *Potentilla fulgens* against liver fluke vector snail *Indoplanorbis exustus*. *Asian J Animal Sci*. 2018b; 12: 30-35.
17. Kumar P. Effect of medicinal plant *Potentilla fulgens* against fecundity, hatchability and survival of *Fasciola* host snail *Indoplanorbis exustus*. *Indian J Scienti Res*. 2021; 11: 19-24.
18. Hajare SW, Lonare MK, Kumar D. Paralytic effect of *Peganum harmala* Linn. Alcoholic extract on *Fasciola gigantica*. *J Vet Pharmacol Toxicol*. 12; 11: 53-55.
19. Kumar P, Singh DK. In vitro anthelmintic activity of *Allium sativum*, *Ferula asafoetida*, *Syzygium aromaticum* and their active components against *Fasciola gigantica*. *J Biol Earth Sci*. 2014; 4: B57-B65.
20. Robertson JL, Russell RM, Preciter HK, Savin NE. Bioassay with arthropods data, 2nd Eds Taylor and Francis, CRC Press. 2007; 1-224.
21. Kumar P, Sunita K, Singh RN, Singh DK. *Fasciola* larvae: Anthelmintic activity of medicinal plant *Potentilla fulgens* against sporocyst, redia and cercaria. *Asian J Adv Res*. 2020; 3: 24-30.
22. Akhar MS, Zafar Iqbal Khan MN, Lateef M. Anthelmintic activity of medicinal plants with particular reference to their use in indo Pakistan sub continent. *Small Rumin*. 2000; 38: 99-107.
23. Kumar P, Sunita K, Singh DK. In vitro activity of different phytochemicals in binary combinations against *Fasciola gigantica*. *Current Life Sci*. 2016; 2: 58-63.
24. Hemalatha S, Laloo D, Prasad SK, Krishnamurthy S. Gastroprotective activity of ethanolic root extract of *Potentilla fulgens* Wall. *Ex Hook. J Ethnopharmacol*. 2013; 146: 505-514. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/23384786/>
25. Roy B, Swargiary A, Syiem D, Tandon V. *Potentilla fulgens* (Family Rosaceae), a medicinal plant of North-East India: A natural anthelmintic. *J Parasit Dis*. 2010; 34: 83-88. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/21966126/>
26. Jaitak V, Kaul VK, Himlata N, Kumar B, Singh J, Dhar, Sharma OP. New hopane triterpenes and antioxidant constituents from *Potentilla fulgens*. *Net Prod Commun*. 2010; 5: 1561-1566. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/21121248/>
27. Hoste H, Jackson F, Athanasiadou S, Thamsborg SM, Hoskin SO. The effects of tannin-rich plants on parasitic nematodes in ruminants. *Trends Parasitol*. 2006; 22: 253-261. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/16632404/>
28. Hoste H, Martinez-Ortiz-De-Montellano C, Manolaraki F, Brunet S, Ojeda-Robertos N, et al. Direct and indirect effects of bioactive tannin-rich tropical and temperate legumes against nematode infections. *Vet Parasitol*. 2012; 186: 18-27. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/22188981/>
29. Athanasiadou S, Kyriazakis I, Jackson F, Coop RL. The effects of condensed tannins supplementation of foods with different protein content on parasitism, food intake and performance of sheep infected with *Trichostrongylus colubriformis*. *Bri J Nutrit*. 2001; 86: 697-706. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/11749679/>
30. Molan AL, Waghorn GC, Mcnabb WC. Effect of condensed tannins on egg hatching and larval development of *Trichostrongylus colubriformis* in vitro. *Vet Rec*. 2002; 150: 65-69. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/11837588/>
31. Rosangkima G, Prasad SB. Antitumor activity of some plants from Meghalaya and Mizoram against Murine ascites Dalton's lymphoma. *Indian J Exp Biol*. 2004; 42: 981-988. **PubMed:** <https://pubmed.ncbi.nlm.nih.gov/15511001/>
32. Majaw S, Challam SK, Syiem D. Effect of *Potentilla fulgens* L. on selective enzyme activities and altered tissue morphology in diabetic mice. *J Morphol Sci*. 2018; 38: 153-160.